

Rangemate 32-10 (B400) Clmrx

49c-3060

Type B Medicated Feed

10517

(For Beef Cattle in Confinement Only)

For increased rate of weight gain and improved feed efficiency in growing-finishing cattle fed in confinement for slaughter. Do not feed undiluted. Do not feed to lactating dairy cattle.

Active Drug Ingredient

Lasalocid (as lasalocid sodium) 400.0 g/ton

Guaranteed Analysis

Crude Protein Min. 32.00 %

This includes no more than 25% equivalent protein from non-protein nitrogen.

Crude Fat Min. 10 %

Crude Fiber Max. 0.1 %

Calcium (Ca) Min. 0.01% Max. 0.50 %

Phosphorus (P) Min. 0.80 %

Salt Min. 3.00% Max. 4.00 %

Cobalt (Co) Min. 4.6 ppm

Copper (Cu) Min. 95 ppm

Iodine (I) Min. 15.0 ppm

Manganese (Mn) Min. 370 ppm

Zinc (Zn) Min. 310 ppm

Selenium (Se) Min. 2.0 ppm

Vitamin A Min. 60,000 IU/lb

Vitamin D Min. 15,000 IU/lb

Vitamin E Min. 90 IU/lb

Dry Matter Min. 50.0 %

Moisture Max. 50.0 %

Ingredients:

Condensed Fermented Corn Extractives, Whey permeate, Urea, Corn Oil, Glycerin, Ammonium Polyphosphate, Salt, Zinc Polysaccharide Complex, Copper Polysaccharide Complex, Manganese Polysaccharide Complex, Ethylenediamine Dihydriodide, Xanthan Gum, Hydrolyzed yeast, Yeast extract, Yeast culture, Coconut oil, Maltodextrin, Lactic Acid, Sodium Selenite, Vitamin A Supplement, Vitamin D₃ Supplement, Vitamin E Supplement, Cobalt Sulfate.

CANNOT BE FED FREE-CHOICE

CAUTION: USE ONLY AS DIRECTED

Manufactured For Rangemate America, LLC

2475 26th Street

Mandan, ND 585545

Liquid Bulk Shipment

Mixing Directions

Recirculate or agitate immediately prior to use for not less than 10 minutes, moving not less than 1% of the tank contents per minute from the bottom of the tank to the top or creating a turbulence at the bottom of the tank that is visible at the top. Recirculate daily, even when not in use.

Directions for Use

Milligrams of Lasalocid per pound = 200.0

Minimum Feeding Rate - lbs/hd/day = 0.25

Maximum Feeding Rate - lbs/hd/day = 2.40

Selenium provided at maximum feeding rate - mg/hd/day = 2.16

which is below the maximum concentration permitted by Federal Regulation Title 21 CFR 573.920. Inclusion of this product in complete feed may need to be reduced if other sources of selenium are utilized.

Caution

The safety of Lasalocid for use in unapproved species has not been established. Do not allow horses or other equines access to feeds containing Lasalocid as ingestion may be fatal. Mix thoroughly with grain and/or roughage prior to feeding. Feeding undiluted or mixing errors may result in an excess Lasalocid concentration which could be fatal to cattle. Inadequate recirculation or agitation may also result in excess Lasalocid concentration fattle to cattle. Medicated liquid supplements stored in recirculating systems: Recirculate immediately prior to use moving no less than 1 percent of the tank contents per minute from the bottom of the tank to the top for not less than 10 minutes. Recirculate daily, even when supplement is not used. If stored in mechanical, air, or other agitation-type tank systems: Agitate immediately prior to use for not less than 10 minutes, creating turbulence at the bottom of the tank that is visible at the top. Agitate daily, even if supplement is not used.

Warning

Do not feed to lactating dairy cattle. A withdrawal time has not been established for pre-ruminating calves. Do not use in calves to be processed for veal.

This product was manufactured in a facility that does not handle or store animal protein products prohibited in ruminant feed as described in 21 CFR section 589.2000.

Expiration Date: _____

8 weeks post manufacture

Because of management decisions and practices beyond our control, seller makes no warranty, expressed or implied, as to description, utility, productiveness, or any other matter, except as to guaranteed analysis.