

## Rangemate 32-10 LI Clmnx

49C-3065

## Directions for Use

*Protein, energy, vitamin and trace mineral supplement for beef and non-lactating dairy cattle*

22017

### Guaranteed Analysis

Crude Protein		Min.	32.00	%
(This includes no more than	25.0	% equivalent protein		
	from non-protein nitrogen)			
Crude Fat		Min.	10.0	%
Crude Fiber		Max.	0.1	%
Calcium (Ca)	Min.	0.01%	Min.	0.50 %
Phosphorus (P)			Min.	0.80 %
Salt (NaCl)	Min.	3.00%	Max.	4.00 %
Cobalt (Co)			Min.	4.6 ppm
Copper (Cu)			Min.	95 ppm
Iodine (I)			Min.	15 ppm
Manganese (Mn)			Min.	370 ppm
Zinc (Zn)			Min.	310 ppm
Selenium			Min.	2 ppm
Vitamin A			Min.	60,000 IU/lb
Vitamin D			Min.	15,000 IU/lb
Vitamin E			Min.	90 IU/lb
Dry Matter		Min.	50.0	%
Moisture		Max.	50.0	%

### Ingredients:

Condensed Fermented Corn Extractives, Whey permeate, Urea, Corn Oil, Glycerin, Hydrochloric acid, Salt, Phosphoric acid, Magnesium chloride, Sulfuric acid, Manganese polysaccharide complex, Zinc polysaccharide complex, Copper polysaccharide complex, Hydrolyzed yeast, Yeast extract, Yeast culture, Coconut oil, Maltodextrin, Lactic Acid, Xanthan gum, Sodium selenite, Cobalt Sulfate, Ethylenediamine Dihydriodide, Vitamin A Supplement, Vitamin D<sub>3</sub> Supplement, Vitamin E Supplement.

**CAUTION: USE ONLY AS DIRECTED** - Use only as recommended on the right side of this page

Manufactured for RangeMate America, LLC

2475 26th Street  
Mandan, ND 58554  
Liquid Bulk Shipment

This supplement is intended for cattle only. Do not feed to sheep. This is a supplement, not a complete ration, and must be fed with adequate roughage and grain.

This supplement may be fed top-dressed, mixed into other components of the ration, or provided in a free-choice lick wheel feeder.

When fed for the first time, or after feeders have been empty, be sure cattle have other feeds available and feed the supplement so that intake is no more than 1.5 lb/hd/day for the first week.

This supplement includes minerals and vitamins, but a free-choice mineral/vitamin supplement is recommended to compensate for variations in ration composition and animal needs. Daily intake should range from 1 to 3 lb/hd/day for a 1000 lb animal. If intake is outside this range, check the following:

**Tank Location** - Intake is higher when tanks are located near water, mineral feeders, shelter, or other areas frequented by cattle.

**Calves** - calves can and should consume the liquid. Their intake must be allowed for also.

**Supplemental feedstuffs** - If cattle are overconsuming, the other feedstuffs may be short of phosphorus or energy. Be sure the mineral includes adequate phosphorus and that the total ration is adequate in energy.

**Caution** - Do not feed to sheep. This product includes supplemental Copper.

Seller makes no warranty, expressed or implied, as to description, utility, productiveness, or any other matter except as to guaranteed analysis.

**This product was manufactured in a facility that does not handle or store animal protein products prohibited in ruminant feed as described in 21 CFR section 589.2000.**